

THE JOURNAL OF JAPANESE BOTANY.

EDITED BY

TOMITARÔ MAKINO, Sc. D.

LECTURER OF BOTANY IN THE FACULTY OF SCIENCE,
IMPERIAL UNIVERSITY OF TOKYO.

Vol. VIII. No. 3.

March 10, 1932.

(Whole No. 73.)

Published by the Tsumura Laboratory, Tokyo, Nippon.

Notes on Japanese Lichens. II.

By

Yasuhiko Asahina.

On the Occurrence of *Catillaria Bouteilli* (DESM.) A. ZAHLBR. in Japan

Recently it was observed several times in the vicinity of Tokyo an epiphyllous lichen growing on coniferous plants, which I could identify with *Catillaria Bouteilli*.

The hosts are :

1. ***Abies firma* SIEB. ET ZUCC.**

Locality Asakawa, Prov. Musashi (leg. A. YAMAMOTO, Sep. 1929).

2. ***Thujopsis dolabrata* SIEB. ET ZUCC.**

Locality Hayamamura, Prov. Sagami (leg. K. HISAUCHI, Nov. 1931).

3. ***Chamaecyparis obtusa* SIEB. ET ZUCC.**

Locality Tomioka, Prov. Musashi (leg. K. HISAUCHI, Jan. 1932).

In the first case the fruits were minute, 0.1-0.2mm across and very scarce but the asci could be observed. Spores oblong, colourless, 1-septate, $8-10 \times 3-4 \mu$ in size. In the latter the fruits were numerous and more developed, 0.2-0.4 mm across.

(Compare the figures in the Japanese text).
