
BRITISH LICHEN SOCIETY BULLETIN No. 77 Winter 1995



NEW, RARE AND INTERESTING BRITISH LICHEN RECORDS

(Contributions to this section are always welcome. Please submit entries to Chris Hitch, The Whin, Wadd Lane, Snape, Saxmundham, Suffolk IP17 1QY, in the form of species, habitat, locality, VC no, VC name, Grid Reference (GR), altitude, where applicable, in metres (m), date, comments and recorder. An authority with date after species in only included, when the record is new to the British Isles. In the interests of accuracy, typecript is much appreciated. Please only use one side of the paper.)

Abrothallus suecicus: on blackened thalli of Ramalina farinacea on Acer pseudoplatanus, Kindrogan Field Centre, Strath Ardle, VC 89, East Perthshire, GR 37/05-62-, 1995. New to Scotland, and apparently a new host.

AM & BJ Coppins

Arthonia zwackhii: on Quercus on N side of River Dart, near junction with River Webburn, Holne Chase, VC 3, South Devon, GR 20/71-71-, 1995.

A M & B J Coppins

Bacidia friesiana: on Sambucus and rusting iron in chalk pit nature reserve, Little Blakenham, VC 25, East Suffolk, GR 62/10-49-, April 1995. New to the County. Determined B J Coppins.

PM Earland-Bennett & CJB Hitch

Bacidia subfuscula: on brick garden wall, Ipswich, VC 25, East Suffolk, GR 62/18-42-, November 1994. Confirmed B J Coppins.

PM Earland-Bennett

Buellia arborea: on lignum of old fence rails with Protoparmelia oleagina [both collections], S of Lower Meads, Rogate, VC 13, West Sussex, GR 41/80-23-, 1970, coll B J Coppins; Shobrooke Park, Crediton, VC 3, South Devon, GR 21/85-01-, 1970, coll B J Coppins & F Rose. New to England. Both collections confirmed by TLC (atranorin and placodiolic acid). B. arborea is easily mistaken in the field for B. griseovirens. The P. oleagina is new to Devon.

B J Coppins

Caloplaca lucifuga: on trunk of ancient Quercus in grounds of Dawyck House, Stobo, VC 78, Peebles, GR 36/16-35-, 1995. New to Scotland.

AM & BJ Coppins

Collema limosum: on garden lawn, Kesgrave, Ipswich, VC 25, East Suffolk, GR 62/23-45-, May 1995.

P M Earland-Bennett

Cornutispora lichenicola: on thallus of Lecanora conizaeoides on twigs of Crataegus bush on top of Roman wall, Colchester, VC 19, North Essex, GR 62/00-25-, January 1994. Determined E Punithalingam.

P M Earland-Bennett

Endococcus parietinus: on thallus of Rinodina gennarii on mortar of old wall at coast, Boyton, VC 25, East Suffolk, GR 62/39-47-, August 1991. Only the second known record on this host. Confirmed D L Hawksworth.

P M Earland-Bennett, C J B Hitch & P N Cayton.

Epicladonia sandstedei: lichenicolous on the thallus of *Cladonia humilis* on gravelly soil, Wick Farm Quarry, VC 25, East Suffolk, GR 62/01-35-, September 1994. New to Suffolk. Confirmed D L Hawksworth.

CJB Hitch

Gelatinopsis ericetorum: on thallus of Baeomyces roseus, North Hill, VC 3, South Devon, GR 31/09-06-, 1991. New to S W England. Determined B J Coppins.

B Benfield

Graphina pauciloculata: on Ilex by East Webburn River, Lizwell Wood, Holne Chase, VC 3, South Devon, GR 20/71-73-. Third major site in SW England.

AM & BJ Coppins

Lecanora navarrensis: on Quercus trunk, 9 km ENE of Ballater, Dinnet Oakwood NNR, VC 92, South Aberdeen, GR 37/4—9—, 1994. New to Scotland.

A M & B J Coppins & G Kantvilas

Lecanora pannonica: on vertical sandstone on the south side of the church, Great Malvern Priory, VC 37, Worcester, GR 32/77-45-, October 1994, and on an east-facing, sandstone window-sill, St Mary's church, Hanley Castle, VC 37, Worcester, GR 32/83-41-, October 1994. New to Worcestershire and much to the west of other recorded sites.

TW Chester

Lecanora pruinosa: on window sills and chamfered plinths on all sides of

the church, on a limestone chest tomb (1855), a coped tomb and a 1914-18 war memorial, Islip churchyard, VC 32, Northampton, GR 42/94-85-, August 1995. New to Northamptonshire and easily the most north-easterly site so far for this recently re-discovered species.

T W Chester

Lecidea lapillicola Vain. (1883): on side of siliceous boulder in woodland, 9 km ENE of Ballater, Dinnet Oakwood NNR, VC 92, South Aberdeen, GR 37/4—9—, 1994. New to the British Isles. This species will be transferred to *Micarea* in a forthcoming paper [some possible earlier names need to be checked]. It is similar to *M.erratica*, but has a thallus of minute granular areoles and a "micareoid" photobiont.

A M & B J Coppins & G Kantvilas

Licea parasitica: growing amongst lichens together with Ramonia interjecta on bark of branches of Sambucus, Baythorne End, VC 19, North Essex, GR 52/72-42-, April 1994. This minute myxomycete may be found growing among or on lichen species but it is not parasitic (B Ing, pers comm). It is the commonest corticolous species of myxomycete in the British Isles. I have found it several times in Essex and Suffolk, although it is more common outside East Anglia. It is most common on Sambucus, but is not confined to this phorophyte. Confirmed E Punithalingam.

PM Earland-Bennett

Macentina abscondita: on shaded Sambucus in chalk pit nature reserve, Little Blakenham, VC 25, East Suffolk, GR 62/10-49-, April 1995. Only the sixth known record of this species, of which three are from East Suffolk!

PM Earland-Bennett & CJ B Hitch

Melaspilea amota: on Quercus with Rinodina isidioides, Stubbs Wood, New Forest, VC 11, South Hampshire, GR 41/3—0—, 1974, collected F Rose & S R Davey; on Quercus on N side of River Dart, Holne Chase, VC 3, South Devon, GR 20/72-72-, 1995, collected B J and A M Coppins - New to England. On Quercus, Talyllyn, Doly-y-cae, W side of Nant Cader, VC 48, Merioneth, GR 23/72-11-, 1972, coll B J Coppins - New to Wales. This species was previously regarded as endemic to southern Ireland. Representative specimens in E.

B J Coppins

Micarea curvata: on side of flat pebble in area of stablilized, riverside shingle, upper reaches of Whiteadder Water, by Johnscleugh, Lammermuir Hills, VC 82, East Lothian, GR 36/63-66-, alt 290m 1995. Second British

record. As with the material from Torrs Warren (Bulletin 74: 60), this too had a K+ violet epithecium. The upper side of the pebble was colonized by Lecidea plana, Porpidia crustulata, P. soredizodes, P. tuberculosa and Rhizocarpon obscuratum.

BJ&AM Coppins

Nectriella robergei: growing with Illosporium carneum on thallus of Peltigera didactyla on soil of old railway sidings, Eight Ash Green, Colchester, VC 19, North Essex, GR 52/95-25-. June 1995.

P M Earland-Bennett & J F Skinner

Opegrapha parasitica: on thallus of Aspicilia calcarea. Selected examples: Dolebury Warren, Mendip Hills, VC 6, North Somerset, GR 31/44-58-, 1984 (E); Port Eynon, Gower, VC 41, Glamorgan, GR 21/46-84-, 1990 (E); Dunkerron, North Kerry, VC H1, 1800s, collected T. Taylor (BM - as 'O. cerebrina'). This species is apparently confined to Aspicilia calcarea, and in the recent British Checklists was provisionally referred to 'O. monspeliensis Nyl.', which is probably a synonym of O. parasitica. The Opegrapha on foveolate pyrenocarpous species of the Verrucariaceae, which has often been called "O. parasitica" (e.g. Flora, p. 411), should be called O. rupestris Pers. (1794).

B J Coppins

Opegrapha rotunda Hafellner (1994) [see "Literature Pertaining" in previous issue]: lichenicolous on thallus of *Physconia distorta*, specimens of mine in E determined from three localities; on tree branch, SW of Cannich, Balcladaich, VC 96, Easterness, GR 28/29-25-, 1975; on *Populus*, between Aberfeldy and Weems, VC 88, Mid-Perth, GR 27/84-49-, 1976; on *Fraxinus*, Dunkroisk, Glen Lochay, VC 88, Mid-Perth, GR 27/5—3—, 1978. New to the British Isles.

B J Coppins

Parmelia conspersa: on slate roofs, Wakes Colne, VC 19, North Essex, GR 52/88-28-, May 1995. This is the first Essex record for thirty years and is present at this site in great quantity.

P M Earland-Bennett & C J B Hitch

Parmelia protomatrae: on slate roof, Framlingham, VC 25, East Suffolk, GR 62/28-61-, May 1995. Third British record - all from East Suffolk! Determined B J Coppins.

PM Earland-Bennett & CJB Hitch

Parmelia pulla: on slate roof, Campsey Ash, VC 25, East Suffolk, GR 62/32-55-, May 1995.

PM Earland-Bennett & CJB Hitch

Phaeosporobolus alpinus R. Sant., Alstrup & D Hawksw, in Alstrup & Hawksworth (1990) [see Bulletin 67: 42]. On thallus of Pertusaria multipuncta, N of Buckland Bridge, Holne Chase, VC 3, South Devon, GR 20/71-72-, alt 100 m. New to the British Isles. Can be distinguished from P. usneae by its smaller conidia (mostly 10-15 vs 15-25 μm) which have smaller cells (3-4 vs 4-6 μm).

AM & BJ Coppins

Plectocarpon sampaianae Diederich & Etayo (1994) [see "Literature Pertaining" in previous issue]: lichenicolous on thallus of Pannaria sampaiana, specimens of mine in E recently identified from two further localities, 3 km E of Drochaid Sgainnir, Glenfinnan, VC 97, Westerness, GR 17/93-79-, 1978; Glasdrum NNR, VC 98, Main Argyll, GR 27/00-45-, 1976.

B J Coppins

Protoparmelia oleagina: - see entry under Buellia arborea (above).

Psammina stipitata: on green coccoid algae on young Fraxinus, Clopton, VC 25, East Suffolk, GR 62/22-54-, May 1995.

PM Earland-Bennett

Pyrenopsis impolita with Placynthium flabellosum on N-facing, steeply sloping, flushed rocks, Traprain Law, VC 82, East Lothian, GR 36/58-74-, alt c. 180 m. New to southern Scotland.

B J & A M Coppins

Sphaerellothecium propinquellum (Nyl.) Rouz & Triebel (1994) [see "Literature Pertaining" in this issue]: in apothecia of Lecanora carpinea on Sorbus, 3 km NW of Tomintoul, Wester Fodderletter, VC 94, Banff, GR 38-14-20-, 1990. New to the British Isles.

BJ&AM Coppins

Steinia geophana: on soil of old railway sidings, Eight Ash Green, Colchester, VC 19, North Essex, GR 52/95-25-, June 1995.

PM Earland-Bennett & JF Skinner

Stigmidium congestum (Körb.) Triebel (1991): in apothecia of Lecanora chlarotera on Populus tremula, 9 km ENE of Ballater, Dinnet Oakwood

NNR, VC 92, South Aberdeen, GR 37/4-9-, 1994. New to the British Isles.

BJ&AM Coppins & G Kantvilas

Usnea fulvoreagens: on Betula, Culbin Forest, VC 95, Moray, GR 28/99-63-. 1983. New to Scotland.

B J Coppins

Verrucaria bryoctona: on sandy ground in old coastal dune system, Aberlady Bay Nature Reserve, VC 82, East Lothian, GR 36/46-82-, 1991. New to Scotland.

B J Coppins & S Ekman

Verrucaria bryoctona: on the ground in land-fill of road junction, near Fleydmore, NE of Forfar, VC 90, Angus, GR 37/48-51-, 1993. Same site as for Aphanopsis coenosa and Leptogium byssinum. Determined BJ Coppins.

R C Munro

Verrucaria bryoctona: on turf of solifluction terrace, near summit of Mam Sodhail, VC 106, East Ross, GR 28/1—2—, alt c1000 m. First montane record.

A M Fryday

Wentiomyces lichenicola ssp. bouteillei: on thallus of Bacidia caligans on mossy top of brick garden wall (10 metres from my front door!), Ipswich, VC 25, East Suffolk, GR 62/18-43-, February 1995. Third British record. Determined D L Hawksworth.

P M Earland-Bennett

Xanthoria elegans: many thalli on concrete tops of walls in urban area, Colchester, VC 19, North Essex, GR 52/95-24-, May 1995. also at Colchester, VC 19, North Essex, GR 62/01-26-, April 1995.

P M Earland-Bennett

Xanthoria elegans: on top of concrete post, Ipswich, VC 25, East Suffolk, GR 62/15-47-, March 1995; also present on concrete tops of walls at Stowupland, VC 25, East Suffolk, GR 62/06-59-, March 1995; Martlesham, VC 25, East Suffolk, GR 62/24-56-, May 1995; Ipswich, VC 25, East Suffolk, GR 62/17-45-, June 1995.

P M Earland-Bennett