
MYCOTAXON

ISSN (print) 0093-4666 (online) 2154-8889 Mycotaxon, Ltd. ©2022

July–September 2022—Volume 137, pp. 545–554

<https://doi.org/10.5248/137.545>

Gender of fungal generic names ending in *–trema*

SHAUN R. PENNYCOOK

*Manaaki Whenua – Landcare Research,
Private Bag 92170, Auckland 1072, New Zealand*
CORRESPONDENCE TO: PennycookS@LandcareResearch.co.nz

ABSTRACT—Fungal generic names ending in *–trema* are reviewed. Most are derived from the latinised Greek neuter noun meaning “perforation; aperture; opening; orifice.” However, some are Latin feminine, with *–trema* referring to a relationship or similarity to *Tremella*.

KEY WORDS—etymology, epithet gender, lichens, nomenclature, *Tremellales*

Introduction

Numerous fungal generic names ending in *–trema* are derived from the Greek neuter noun τρῆμα, meaning “perforation; aperture; opening; orifice.” Most of these are lichen genera, with names referring to characteristics of their ascromatal ostioles.

It has been widely assumed that all generic names ending in *–trema* have neuter gender, regardless of their etymology. However, there are several tremellalean genera where the ending *–trema* has been treated as a Latin feminine, apparently intended as an abbreviation alluding to their affinity with *Tremella*.

A similar mix of implied genders occurs among botanical and algal names. Of 14 plant genera ending in *–trema* (IPNI 2022), nine have neuter species epithets, four have feminine epithets, and one has no listed species. Of seven algal genera ending in *–trema* (AlgaeBase 2022), two have neuter species epithets, three have feminine epithets, and two have no listed species; moreover the AlgaeBase website explicitly lists the “accepted gender” of three of these genera—one as neuter and two as feminine. The protoglosses of these botanical

and algal genera need to be checked to confirm whether or not they are derived from Greek τρῆμα.

Here, a listing of all fungal *-trema* genera is presented, with the evidence for their gender compiled and annotated, including comments on the orthography of higher taxon names typified by such genera.

Materials & methods

Two of the three major fungal nomenclature websites, Index Fungorum (2022) and MycoBank (2022), were interrogated to compile a list of all fungal *-trema* genera. Evidence of their gender was ascertained by investigating: (a) the gender of adjectival epithets published by the original and subsequent authors of species; (b) etymologies (either published or implied by the generic descriptions); and (c) the orthography of higher taxa typified by them.

Results & discussion

Fifty fungal genera ending with *-trema* were located and classified into four groups: 35 lichenised neuter genera; eight non-lichenised neuter genera; six Latin feminine genera; and one microsporidial (feminine?; neuter?) genus.

Lichenised genera with names based on Greek τρῆμα (neuter)

Mycasterotrema is a nomen nudum; and *Prototrema* is rather obscure and apparently has no named species. All but two of the remaining 33 genera in this group have neuter adjectival epithets, either in the holotype or in other species (if the holotype epithet is gender uninformative). *Sanguinotrema* (*Graphidaceae*) is a monotypic genus with an uninformative epithet. *Xerotrema* (*Odontotremataceae*) was published as a monotypic genus with a feminine holotype epithet and a subsequently included second species has an uninformative epithet; I have corrected the holotype epithet to the neuter form.

Four tribes, four families, and one order are typified by genera in this group; these names are formed from Greek neuter linguistic stems, resulting in the terminations *-tremataeae*, *-tremataceae*, or *-trematales*.

Acanthotrema Frisch (Frisch 2006: 77)

Type: *Acanthotrema brasiliannum* (Hale) Frisch

Amazonotrema Kalb & Lücking (Kalb 2009: 18)

Type: *Amazonotrema nigrum* Kalb & Lücking

Ampliotrema Kalb (Frisch 2006: 81)

Type: *Ampliotrema amplius* (Nyl.) Kalb [a neuter comparative epithet]

Asterotrema Müll. Arg. (Müller [Argovensis] 1884: 19)

Type: *Asterotrema parasiticum* Müll. Arg.

Austrotrema I. Medeiros, Lücking & Lumbsch (Medeiros & al. 2017: 14)

Type: *Austrotrema bicinctulum* (Nyl.) I. Medeiros & al.

Borinquenotrema Merc.-Díaz, Lücking & Parnmen (Mercado-Díaz & al. 2014: 191)

Type: *Borinquenotrema soredicarpum* Merc.-Díaz & al.

Byssotrema M. Cáceres, Aptroot & Lücking (Cáceres & al. 2014: 91)

Type: *Byssotrema mirabile* M. Cáceres & al.

Clandestinotrema Rivas Plata, Lücking & Lumbsch (Rivas Plata & al. 2012a: 116)

Type: *Clandestinotrema clandestinum* (Ach.) Rivas Plata & al.

Coccotrema Müll. Arg. (Müller [Argovensis] 1889: 171)

Type: *Coccotrema antarcticum* Müll. Arg.

Coccotremataceae Henssen ex J.C. David & D. Hawksw. (David & Hawksworth 1991: 14)

Compositrema Rivas Plata, Lücking & Lumbsch (Rivas Plata & al. 2012b: 1172)

Type: *Compositrema cerebriforme* J.E. Hern. & Lücking

Conotrema Tuck. (Tuckerman 1848: 278)

Type: *Conotrema urceolatum* (Ach.) Tuck.

Cruentotrema Rivas Plata, Papong, Lumbsch & Lücking (Rivas Plata & al. 2012a: 116)

Type: *Cruentotrema cruentatum* (Mont.) Rivas Plata & al.

Cryptoschizotrema Aptroot, Lücking & M. Cáceres (Hyde & al. 2019: 129)

Type: *Cryptoschizotrema cryptotrema* (Nyl.) Aptroot & al.

Additional sp.: *Cryptoschizotrema minus* E.L. Lima & Lücking [a neuter comparative epithet] (Lima & al. 2019: 417)

Enigmotrema Lücking (Sipman & al. 2012: 66)

Type: *Enigmotrema rubrum* Lücking

Glaucotrema Rivas Plata & Lumbsch (Rivas Plata & al. 2012b: 1174)

Type: *Glaucotrema glaucophaeum* (Kremp.) Rivas Plata & Lumbsch

Gymnotrema Nyl. (Nylander 1858: 119)

Type: *Gymnotrema atratum* (Fée) Nyl.

Gyrotrema Frisch (Frisch & Kalb 2006: 379)

Type: *Gyrotrema sinuosum* (Sipman) Frisch

Leptotrema Mont. & Bosch (Montagne 1856: 363)

Syntypes: *Leptotrema zollingeri* Mont. & Bosch; *L. prevostianum* (Mont.) Mont.

Leptotremataceae Lumbsch, Kraichak & Lücking (Lumbsch & al. 2014: 47)

Melanotrema Frisch (Frisch & Kalb 2006: 382)

Type: *Melanotrema platystomum* (Mont.) Frisch

Mycasterotrema Räsänen (Räsänen 1943: 22, as “[435. Mycasterotrema Räs.”], nom. nud.)

= *Asterotrema* Müll. Arg. (Lamb 1963: 421)

Myriotrema Fée (Fée 1825: XLIX, 103)

Type: *Myriotrema olivaceum* Fée

Odontotrema Nyl. (Nylander 1858: 143)

Type: *Odontotrema phacidiooides* Nyl.

Additional sp.: *Odontotrema minus* Nyl. [a neuter comparative epithet] (Nylander & Saelan 1859: 91)

Odontotremataceae D. Hawksw. & Sherwood (Hawksworth & Sherwood 1982: 263)

Odontotrematales Lücking (Lücking 2019: 233)

Parmotrema A. Massal. (Massalongo 1860: 248)

Type: *Parmotrema perforatum* (Jacq.) A. Massal.

Phaeotrema Müll. Arg. (Müller [Argovensis] 1887: 10)

Type: *Phaeotrema subfarinosum* (Fée) Müll. Arg.

Plagiotrema Müll. Arg. (Müller [Argovensis] 1885: 387)

Type: *Plagiotrema cubanum* Müll. Arg.

Pleurotrema Müll. Arg. (Müller [Argovensis] 1885: 388)

Type: *Pleurotrema polysenum* (Nyl.) Müll. Arg.

Pleurotremataceae Walt. Watson (Watson 1929: 113)

Prototrema M. Choisy (Choisy 1928: tab. 18)

= *Thelotrema* Ach. (Lamb 1963: 595)

Pycnotrema Rivas Plata & Lücking (Rivas Plata & al. 2012a: 120)

Type: *Pycnotrema pycnoporellum* (Nyl.) Rivas Plata & Lücking

Sanguinotrema Lücking (Lücking & al. 2015: 441)

Type: *Sanguinotrema wightii* (Taylor) Lücking

Sanguinotremataceae Lücking, Kraichak & Lumbsch (Lücking & al. 2015: 442)

Schizotrema Mangold & Lumbsch (Mangold & al. 2009: 348, 657)

Type: *Schizotrema zebrinum* Mangold

Thelotrema Ach. (Acharius 1803: 130)

Type: *Thelotrema lepadinum* (Ach.) Ach.

Thelotremataceae Rivas Plata, Lücking & Lumbsch (Rivas Plata & al. 2012a: 114)

Thelotremataceae Stizenb. [as “*Thelotremae*”] (Stizenberger 1862: 167)

Trichotrema Clem. (Clements 1909: 41, 173)

Type: *Trichotrema trichosporum* (Müll. Arg.) Clem.

Trinathotrema Lücking, Rivas Plata & Mangold (Lücking & al. 2011: 195)

Type: *Trinathotrema stictideum* (Nyl.) Lücking & al.

Wirthiotrema Rivas Plata, Kalb, Frisch & Lumbsch (Rivas Plata & al. 2010: 198)

Type: *Wirthiotrema glaucopallens* (Nyl.) Rivas Plata & Kalb

Additional sp.: *Wirthiotrema duplomarginatum* Lücking & al. (Sipman & al. 2012: 202)

Wirthiotremataceae Lumbsch, Kraichak & Lücking (Lumbsch & al. 2014: 47)

Xerotrema Sherwood & Coppins (Sherwood & Coppins 1980: 368)

Type: *Xerotrema megalosporum* Sherwood & Coppins [as "megalospora"]

Non-lichenised genera with names based on Greek τρῆμα (neuter).

The eight genera in this group are from diverse asco- and basidiomycete families or orders (as annotated below). *Hydnotrema* is illegitimate and *Sistotrema* Pers. is an unavailable earlier homonym of a sanctioned name; both have been synonymised with other legitimate genera. The other six genera have either neuter adjectival or gender uninformative epithets. In some of these generic names, latinised Greek neuter *trema* has been used as an ending analogous to latinised Greek neuter *stoma* (with very similar meanings); e.g., *Lophiotrema* was proposed as a sister genus to *Lophiostoma* (Saccardo 1878: 338–339). Three of the other generic names were created by adding prefixes to *Lophiotrema*.

Four families are typified by genera in this group; these names are formed from Greek neuter linguistic stems, resulting in the termination *-tremataceae*.

Antealophiotrema A. Hashim. & Kaz. Tanaka (Hashimoto & al. 2017: 68)

[*Pleosporales*]

Type: *Antealophiotrema brunneosporum* (Ying Zhang & al.) A. Hashim. & Kaz. Tanaka

Echinotrema Park.-Rhodes (Parker-Rhodes 1955: 367) [*Hydnodontaceae*]

Type: *Echinotrema clavulare* Park.-Rhodes

Echinotremataceae Jülich (Jülich 1982: 366)

Hydnotrema Link (Link 1833: 298) [*Hydnaceae*], nom. illeg.; ≡ *Sistotrema* Fr. 1821, nom. sanct.

Type: *Hydnotrema confluens* (Pers.) Link; ≡ *Sistotrema confluens* Pers. 1794

Lophiotrema Sacc. (Saccardo 1878: 338) [*Lophiotremataceae*]

Type: *Lophiotrema nucula* (Fr.) Sacc. [a noun in apposition]

Additional spp.: *Lophiotrema alpigenum* (Fuckel) Sacc.; *L. nuculinum* (Rehm)

Sacc.; *L. semilibерум* (Desm.) Sacc.; *L. sexnucleatum* (Cooke) Sacc.

Lophiotremataceae K. Hiray. & Kaz. Tanaka (Hirayama & Tanaka 2011: 405)

Neolophiotrema G.C. Ren & K.D. Hyde (Ren & al. 2021: 28) [*Anteagloniaceae*]

Type: *Neolophiotrema xiaokongense* G.C. Ren & K.D. Hyde

Pseudolophiotrema A. Hashim. & Kaz. Tanaka (Hashimoto & al. 2017: 70)

[*Pseudolophiotremataceae*]

Type: *Pseudolophiotrema elymicola* A. Hashim. & Kaz. Tanaka

Pseudolophiotremataceae K.D. Hyde & Hongsanan (Hongsanan & al. 2018: 97)

Sistotrema Fr. (Fries 1821: 426, nom. sanct.) [*Hydnaceae*]

Type: *Sistotrema confluens* Pers. 1794, nom. sanct.

Sistotremataceae Jülich (Jülich 1982: 390)

Sistotrema Pers. (Persoon 1794: 108) [*Cerrenaceae*]; legitimate but unavailable (ICN

2018: Art. F.3.4); \equiv *Cerrena* Gray (Gray 1821: 649)

Type: *Sistotrema cinereum* Pers.; = *Cerrena unicolor* (Bull.) Murrill (Murrill 1903: 91)

Generic names formed from Latin *trema* (feminine)

Five genera in this group are in *Tremellales* and have species with feminine adjectival epithets. A sixth, *Sclerotrema*, is a monotypic genus in *Auriculariales* but has an etymology explicitly defining *trema* as a reference to tremellaceous fungi; I have corrected its neuter holotype epithet to the feminine form. Two families are typified by genera in this group; these names are formed from Latin feminine linguistic stems, resulting in the termination *-tremaceae*.

Cuniculitrema J.P. Samp. & R. Kirschner (Kirschner & al. 2001: 155) [*Tremellales*]

Type: *Cuniculitrema polymorpha* R. Kirschner & J.P. Samp.

Cuniculitremaceae J.P. Samp., R. Kirschner & M. Weiss (Kirschner & al. 2001: 155)

Gelidatrema Yurkov, Xin Zhan Liu, F.Y. Bai, M. Groenew. & Boekhout (Liu & al. 2015: 138) [*Tremellales*]

Type: *Gelidatrema spencermartiniae* (V. de García & al.) Yurkov & al.

Additional sp.: *Gelidatrema psychrophila* M. Tsuji (Tsuji & al. 2018: 69)

Papiliotrema J.P. Samp., M. Weiss & R. Bauer (Sampaio & al. 2002: 875) [*Tremellales*]

Type: *Papiliotrema bandonii* J.P. Samp. & al.

Additional sp.: *Papiliotrema aurea* (Saito) Xin Zhan Liu & al. (Liu & al. 2015: 126)

Rhynchogastrema B. Metzler & Oberw. (Metzler & al. 1989: 281) [*Tremellales*]

Type: *Rhynchogastrema coronata* B. Metzler & Oberw.

“Etym.: Rhynchos (Gr.) – beak, gaster (Gr.) – body, belly, *Tremella*.”

Rhynchogastremaceae Oberw. & B. Metzler (Metzler & al. 1989: 283)

“*Rhynchogastremataceae*” (Liu & al. 2015: 125) [an orthographic error]

Sclerotrema Spirin & Malysheva (Malysheva & Spirin 2017: 712) [*Auriculariales*]

Type: *Sclerotrema griseobrunnea* (K. Wells & Raity.) Spirin & Malysheva [as “*griseobrunneum*”]

“Etymology. ‘skleros’ (Greek, adj.) – dry, and ‘trema’ – a reference to tremellaceous fungi, . . .”

Sirotrema Bandoni (Bandoni 1986: 668) [*Tremellaceae*]

Type: *Sirotrema pusilla* Bandoni

Genus name in *Microsporidia* (feminine?; neuter?)

Although *Microsporidia* are now categorised as fungi, their names continue to be governed by the International Code of Zoological Nomenclature (ICN 2018: Art. F.1.1).

The etymology of *Systemostrema* identifies the first element *systemos-* as Greek, and the assumption would be that both elements are Greek. However, from the etymology, the authors appear to interpret *-trema* as meaning “thread” (= Greek neuter *nema*). There is nothing in the morphology of the genus that suggests the Greek etymology *-trema*, meaning “opening”.

Like the Botanical Code, the Zoological Code requires adjectival epithets to agree with the gender of the genus (ICZN 2000: Art. 31.2). Two of the five *Systemostrema* species have feminine epithets (the other three are gender uninformative), but there is no connection with *Tremella*. The speculation arises that the authors were either treating *-trema* as a “meaningless” feminine suffix (cf. *-ella* and *-opsis*); or that *-trema* was an error for *-nema*, both of which would have required neuter adjectival epithets.

Systemostrema E.I. Hazard & Oldacre (Hazard & Oldacre 1976: 87) [*Microsporidia*]

Type: *Systemostrema tabani* E.I. Hazard & Oldacre

ETYMOLOGY: “We name this genus *Systemostrema*, meaning “thread tapering to a point” and relating to the polar filament which abruptly constricts to a narrow distal portion.”

Additional spp.: *Systemostrema alba* J.I.R. Larss.; *S. candida* J.I.R. Larss. (Larsson 1988: 16)

Acknowledgments

I thank John David (Royal Horticultural Society, Wisley, UK) for his advice on the etymology of *Systemostrema* and for refereeing the manuscript; Paul Kirk (Royal Botanic Gardens, Kew, Richmond, UK) for arousing my interest in this topic and for refereeing the manuscript; and Tom May (Royal Botanic Gardens Victoria, Melbourne, Australia) for refereeing the manuscript.

Literature cited

- Acharius E. 1803. Methodus que omnes detectus lichenes. Stockholm, F.D.D. Ulrich. 152 p.
- AlgaeBase. 2022. Listing the world's algae. <https://www.algaebase.org/search/> [Accessed: September 2022].
- Bandoni RJ. 1986. *Sirotrema*: a new genus in the *Tremellaceae*. Canadian Journal of Botany 64(3): 668–676. <https://doi.org/10.1139/b86-085>
- Cáceres MES, Aptroot A, Parnmen S, Lücking R. 2014. Remarkable diversity of the lichen family *Graphidaceae* in the Amazon rain forest of Rondônia, Brazil. Phytotaxa 189(1): 87–136. <https://doi.org/10.11646/phytotaxa.189.1.8>

- Choisy M. 1928. *Icones lichenum universalis*, fasc. 3: tab. 17–24.
- Clements FE. 1909. The genera of fungi. Minneapolis, H.W. Wilson Company. 227 p.
- David JC, Hawksworth DL. 1991. Validation of six families of lichenized ascomycetes. *Systema Ascomycetum* 10(1): 13–18.
- Fée ALA. 1825 [“1824”]. Essai sur les crytogames des écorces exotiques officinales, première partie. CII +167 p.
- Fries EM. 1821 *Systema mycologicum* 1. Lund, Berling. 520 p.
- Frisch A. 2006. Contributions towards a new systematics of the lichen family *Thelotremaeaceae*. I. The lichen family *Thelotremaeaceae* in Africa. *Bibliotheca Lichenologica* 92: 3–370.
- Frisch A, Kalb K. 2006. Contributions towards a new systematics of the lichen family *Thelotremaeaceae*. II. A monograph of *Thelotremaeaceae* with a complex structure of the columella. *Bibliotheca Lichenologica* 92: 371–516.
- Gray SF. 1821. Natural arrangement of British plants, vol. 1. London, Baldwin, Craddock, & Joy. 824 p.
- Hashimoto A, Matsumura M, Hirayama K, Tanaka K. 2017. Revision of *Lophiotremataceae* sensu lato (*Pleosporales, Dothideomycetes*): establishment of *Aquasubmersaceae*, *Cryptocoryneaceae*, and *Hermatomycetaceae* fam. nov. *Persoonia* 39: 51–73. <https://doi.org/10.3767/persoonia.2017.39.03>
- Hawksworth DL, Sherwood MA. 1982. Two new families in the *Ascomycotina*. *Mycotaxon* 16(1): 262–264.
- Hazard EI, Oldacre SW. 1976. Revision of *Microsporida* (*Protozoa*) close to *Thelohania*, with descriptions of one new family, eight new genera, and thirteen new species. Technical Bulletin, United States Department of Agriculture 1530. 104 p.
- Hirayama K, Tanaka K. 2011. Taxonomic revision of *Lophiostoma* and *Lophiotrema* based on reevaluation of morphological characters and molecular analyses. *Mycoscience* 52(6): 401–412. <https://doi.org/10.1007/S10267-011-0126-3>
- Hongsanan S, Zeng YJ, Hyde KD, Xie N. 2018. Familial status of *Lophiotremataceae* and its related families in *Pleosporales*. *Phytotaxa* 383(1): 93–102. <https://doi.org/10.11646/phytotaxa.383.1.5>
- Hyde KD, Tennakoon DS, Jeewon R, Bhat DJ, Maharachchikumbura SSN, Rossi W & al. 2019. Fungal diversity notes 1036–1150: taxonomic and phylogenetic contributions on genera and species of fungal taxa. *Fungal Diversity* 96. 242 p.
- ICN. 2018. International Code of Nomenclature for algae, fungi, and plants. *Regnum Vegetabile* 159. 254 p. <https://www.iapt-taxon.org/nomen/main.php?page=pf>
- ICZN. 2000. International Code of Zoological Nomenclature, 4th ed, online. <https://www.iczn.org/the-code/the-code-online/>
- Index Fungorum. 2022. Names search. <http://www.indexfungorum.org/names/Names.asp> [Accessed: January 2022].
- IPNI. 2022. International Plant Names Index (IPNI). <https://www.ipni.org/> [Accessed: September 2022].
- Jülich W. 1982 [“1981”]. Higher taxa of *Basidiomycetes*. *Bibliotheca Mycologica* 85. 485 p.
- Kalb K. 2009. New taxa and new records of thelotremoid *Graphidaceae*. *Herzogia* 22: 17–42.
- Kirschner R, Sampaio JP, Gadanho M, Weiß M, Oberwinkler F. 2001. *Cuniculitrema polymorpha* (*Tremellales*, gen. nov. and sp. nov.), a heterobasidiomycete vectored by bark beetles, which is the teleomorph of *Sterigmatosporidium polymorphum*. *Antonie van Leeuwenhoek* 80(2): 149–161. <https://doi.org/10.1023/a:1012275204498>
- Lamb IM. 1963. Index nominum lichenum inter annos 1932 et 1960 divulgatorum. xii + 810 p.

- Larsson JIR. 1988. On the taxonomy of the genus *Systemostrema* Hazard & Oldacre, 1975 (*Microspora, Thelohaniidae*), with description of two new species. Systematic Parasitology 11: 3–17. <https://doi.org/10.1007/BF00006972>
- Lima EL de, Maia LC, Martins MCB, Silva NL, Lücking R, Cáceres MED. 2019. Five new species of *Graphidaceae* from the Brazilian Northeast, with notes on *Diorygma alagoense*. Bryologist 122(3): 414–422. <https://doi.org/10.1639/0007-2745-122.3.414>
- Link JHF. 1833. Handbuch. Dritter Theil. Berlin, S.J. Joscephy. 536 p.
- Liu XZ, Wang QM, Göker M, Groenewald M, Kachalkin AV, Lumbsch HT, Millanes AM, Wedin M, Yurkov AM, Boekhout T, Bai FY. 2015. Towards an integrated phylogenetic classification of the *Tremellomycetes*. Studies in Mycology 81: 85–147. <https://doi.org/10.1016/j.simyco.2015.12.001>
- Lücking R. 2019. Stop the abuse of time! Strict temporal banding is not the future of rank-based classifications in fungi (including lichens) and other organisms. Critical Reviews in Plant Sciences 38(3): 199–253. <https://doi.org/10.1080/07352689.2019.1650517>
- Lücking R, Rivas Plata E, Mangold A, Sipman HJM, Aptroot A, Gonzalez RM, Kalb K, Chaves JL, Ventura N, Esquivel RE. 2011. Natural history of Nash's pore lichens, *Trinathotrema* (*Ascomycota: Ostropales: Stictidaceae*). Bibliotheca Lichenologica 106: 187–210.
- Lücking R, Mangold AS, Rivas Plata E, Parnmen S, Kraichak E, Lumbsch HT. 2015. Morphology-based phylogenetic binning to assess a taxonomic challenge: a case study in *Graphidaceae* (*Ascomycota*) requires a new generic name for the widespread *Leptotrema wightii*. Botanical Journal of the Linnean Society 179(3): 436–443. <https://doi.org/10.1111/boj.12327>
- Lumbsch HT, Kraichak E, Parnmen S, Rivas Plata E, Aptroot A, Caceres MES & al. 2014. New higher taxa in the lichen family *Graphidaceae* (lichenized *Ascomycota: Ostropales*) based on a three-gene skeleton phylogeny. Phytotaxa 189: 39–51. <https://doi.org/10.11646/phytotaxa.189.1.5>
- Malysheva V, Spirin V. 2017. Taxonomy and phylogeny of the *Auriculariales* (*Agaricomycetes, Basidiomycota*) with stereoid basidiocarps. Fungal Biology 121(8): 689–715. <https://doi.org/10.1016/j.funbio.2017.05.001>
- Mangold A, Elix JA, Lumbsch HT. 2009. *Thelotremae*. 195–406, 653–659, in: PM McCarthy, B Kuchlmayr (eds). Flora of Australia, vol. 57, lichens 5. Melbourne, CSIRO Publishing.
- Massalongo AB. 1860. Esame comparativo di alcune genere di licheni. Atti del Istituto Veneto di Scienze, Lettere ed Arti, Ser. 3, 5: 247–276.
- Medeiros ID, Kraichak E, Lücking R, Mangold A, Lumbsch HT. 2017. Assembling a taxonomic monograph of tribe *Wirthiotremateae* (lichenized *Ascomycota: Ostropales: Graphidaceae*). Fieldiana, Life and Earth Sciences 9(1): 1–31. <https://doi.org/10.3158/2158-5520-16.9.1>
- Mercado-Díaz JA, Lücking R, Parnmen S. 2014. Two new genera and twelve new species of *Graphidaceae* from Puerto Rico: a case for higher endemism of lichenized fungi in islands of the Caribbean? Phytotaxa 189(1): 186–203. <https://doi.org/10.11646/phytotaxa.189.1.14>
- Metzler B, Oberwinkler F, Petzold H. 1989. *Rhynchogastrema* gen. nov. and *Rhynchogastremaceae* fam. nov. Systematic and Applied Microbiology 12(3): 280–287. [https://doi.org/10.1016/s0723-2020\(89\)80075-x](https://doi.org/10.1016/s0723-2020(89)80075-x)
- Montagne JPFC. 1856. Sylloge generum specierumque crytogrammarum. Pais, Baillière. 498 p.
- Müller [Argoviensis] J. 1884. Enumerationis lichenum aegyptiacorum, supplementum primum. Revue Mycologique 6: 15–20.
- Müller [Argoviensis] J. 1885. *Pyrenocarpea* cubenses a cl. D. Wright lectae. Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie 6: 374–421.
- Müller [Argoviensis] J. 1887. *Graphideae* Feeanae. Mémoires de la Société de Physique et d'Histoire Naturelle de Genève 29(8): 1–80.
- Müller [Argoviensis] J. 1889. Lichens. Mission Scientifique du Cap Horn, Tome 5, Botanique: 141–172.

- Murrill WA. 1903. A historical review of the genera of the *Polyporaceae*. *Journal of Mycology* 9(2): 87–102. <https://doi.org/10.2307/3752511>
- MycoBank. 2022. Basic search. <https://www.mycobank.org/page/Basic%20names%20search> [Accessed: January 2022].
- Nylander W. 1858. Énumération générale de lichens, avec l'indication sommaire de leur distribution géographique. *Mémoires de la Société Impériale des Sciences Naturelles de Cherbourg* 5: 85–146.
- Nylander W, Saelan AT. 1859. Herbarium musei fenneci. *Helsingfors, Finska Litteratur-Sällskapets Tryckeri*. 118 p.
- Parker-Rhodes AF. 1955. The basidiomycetes of Skokholm Island. XIII. *Echinotrema clanculare* gen. et sp.nov. *Transactions of the British Mycological Society* 38(4): 366–368. [https://doi.org/10.1016/s0007-1536\(55\)80039-3](https://doi.org/10.1016/s0007-1536(55)80039-3)
- Persoon CH. 1794. Dispositio methodica fungorum. *Neues Magazin für die Botanik* 1: 81–128.
- Räsänen V. 1943. Das System der Flechten. *Acta Botanica Fennica* 33. 82 p.
- Ren GC, Wanasinghe DS, Monkai J, Hyde KD, Mortimer PE, Xu JC, Pang A, Gui H. 2021. Introduction of *Neolophiotrema xiaokongense* gen. et sp. nov. to the poorly represented *Anteagloniaceae* (*Pleosporales, Dothideomycetes*). *Phytotaxa* 482(1): 25–35. <https://doi.org/10.11646/phytotaxa.482.1.3>
- Rivas Plata E, Kalb K, Frisch A. 2010. *Wirthiotrema*: a new genus for the *Thelotrema glaucopallens* group (*Ascomycota: Ostropales: Thelotremonoid Graphidaceae*). *Lichenologist* 42(2): 197–202. <https://doi.org/10.1017/S0024282909990454>
- Rivas Plata E, Lücking R, Lumbsch HT. 2012a. A new classification for the lichen family *Graphidaceae* (*Ascomycota: Lecanoromycetes: Ostropales*). *Fungal Diversity* 52: 107–121. <https://doi.org/10.1007/s13225-011-0135-8>
- Rivas Plata E, Lücking R, Lumbsch T. 2012b. Molecular phylogeny and systematics of the *Ocellularia* clade (*Ascomycota: Ostropales: Graphidaceae*). *Taxon* 61: 1161–1172. <https://doi.org/10.1002/tax.616001>
- Saccardo PA. 1878. *Fungi italici autographice delineati a prof. P.A. Saccardo*. *Michelia* 1(3): 326–350.
- Sampaio JP, Weiß M, Gadango M, Bauer R. 2002. New taxa in the *Tremellales*: *Bulleribasidium oberjochense* gen. et sp. nov., *Papiliotrema bandonii* gen. et sp. nov. and *Fibulobasidium murrhardtense* sp. nov. *Mycologia* 94(5): 873–887. <https://doi.org/10.2307/3761703>
- Sherwood MA, Coppins BJ. 1980. *Xerotrema*, a new genus of ondontotremoid fungus from Scotland. *Notes from the Royal Botanical Garden Edinburgh* 38(2): 367–371.
- Sipman HJM, Lücking R, Aptroot A, Chaves JL, Kalb K, Umaña Tenorio L. 2012. A first assessment of the TICOLICHEN biodiversity inventory in Costa Rica and adjacent areas: the thelotremonoid *Graphidaceae* (*Ascomycota: Ostropales*). *Phytotaxa* 55. 214 p. <https://doi.org/10.11646/phytotaxa.55.1.1>
- Stizenberger E. 1862. Beitrag zur Flechtensystematik. Bericht über die Thätigkeit der St. Gallischen Naturwissenschaftlichen Gesellschaft 1861–62: 124–182.
- Tsuji M, Tanabe Y, Vincent WF, Uchida M. 2018. *Gelidatrema psychrophila* sp. nov., a novel yeast species isolated from an ice island in the Canadian High Arctic. *Mycoscience* 59(1): 67–70. <https://doi.org/10.1016/j.myc.2017.08.010>
- Tuckerman E. 1848. A synopsis of the lichens of the northern United States and British America. *Proceedings of the American Academy of Arts and Sciences* 1: 195–285.
- Watson W. 1929. The classification of lichens. Part II. *New Phytologist* 28: 85–116.