

doi.org/10.3114/fuse.2023.11.01

Cylindromonium dirinariae sp. nov. (Ascomycota, Hypocreales), a new nectrioid lichenicolous species on *Dirinaria applanata* in Japan

A. Ohmaki¹, I. Okane^{2*}, P.W. Crous³, G.J.M. Verkley³

¹Agro-Bioresources Science and Technology, Univ. of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8577, Japan ²Faculty of Life and Environmental Sciences, Univ. of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8577, Japan ³Westerdijk Fungal Biodiversity Institute, Uppsalalaan 8, 3584 CT Utrecht, Netherlands

*Corresponding author: okane.izumi.fw@u.tsukuba.ac.jp

Key words:	Abstract: A nectrioid fungus forming a pinkish colony with mainly solitary phialides producing ellipsoid, aseptate conidia
culture	in mucoid packets was isolated from Dirinaria applanata. Our taxonomic study based on morphology and phylogenetic
inoculation	analysis using ITS rDNA sequences revealed that the isolates represented a member of the genus Cylindromonium. Based
lichenicolous fungi	on further morphological examination, nucleotide sequence comparison, and phylogenetic analysis based on LSU rDNA,
Nectriaceae	tef1, and rpb2 in addition to the phylogenetic analysis using the ITS rDNA sequences, the fungus from Dirinaria represents
new taxon	a new species, which is described here as Cylindromonium dirinariae sp. nov. Furthermore, inoculation experiments
phylogeny	revealed that this species can also produce perithecia when inoculated on the host lichen in laboratory environments.

Citation: Ohmaki A, Okane I, Crous PW, Verkley GJM (2023). Cylindromonium dirinariae sp. nov. (Ascomycota, Hypocreales), a new nectrioid lichenicolous species on Dirinaria applanata in Japan. Fungal Systematics and Evolution **11**: 1–10. doi: 10.3114/fuse.2023.11.01 Received: 8 November 2022; Accepted: 2 January 2023; Effectively published online: 23 January 2023 Corresponding editor: A.J.L. Phillips

INTRODUCTION

Lichenicolous fungi is a term used to circumscribe fungi that grow on lichens. They can interact with their lichen hosts as saprophytes, parasites and commensalistic parasymbionts. Lichenicolous fungi usually establish a symbiotic relationship with a single species or genus of lichens, while some species have a wide host range (Diederich et al. 2018). Approximately 2 300 species of lichenicolous fungi have been described on lichens globally and they are classified into ca. 400 genera, ca. 100 families, 55 orders and 10 classes. Ninety-six percent of the total number of lichenicolous fungi are ascomycetes and four percent of the fungi are basidiomycetes (Diederich et al. 2018). About 166 lichenicolous fungi have been reported from Japan to date (Frisch et al. 2018, Ohmura & Kashiwadani 2018, Tadome et al. 2018, Zhurbenko & Ohmura 2018a, b, Zhurbenko et al. 2018, Zhurbenko & Ohmura 2019, 2020, Frisch et al. 2020, Tadome & Ohmura 2021, 2022, Tadome et al. 2022). In spite of these reports, there has been relatively little research conducted on this fungal group in Japan. Therefore, many species remain to be discovered and described.

The genus *Cylindromonium*, with the type species *C. eugeniicola*, was segregated from *Acremonium* based on analyses of ITS and LSU rDNA sequence data (Summerbell *et al.* 2011, Crous *et al.* 2019b). *Cylindromonium* species are known to be lichenicolous, mycophilic, or saprophytic (Gams 1971, Crous *et al.* 2019b, 2020, 2021). *Cylindromonium* was established as a

genus to accommodate acremonium-like taxa with unbranched, hyaline, phialidic conidiophores, and cylindrical 1-septate conidia (Crous *et al.* 2019b). A total of five asexual species have been assigned to *Cylindromonium*, for which no sexual morph has thus far been reported (Crous *et al.* 2019b, 2020, 2021). During our research on the diversity of lichenicolous fungi in Japan, a fungus colonising the *Physciaceae* lichen *Dirinaria applanata* was found. The purpose of this study is to describe the morphological, physiological, and ecological features of this species, clarify the link to its sexual morph, and discuss its taxonomic placement.

MATERIALS AND METHODS

Collection materials

Field investigations were performed from September 2020 to March 2021 in Tsukuba city, Ibaraki prefecture, Japan. Specimens of the fungus growing on the lichen host *D. applanata* were found on the bark of *Zelkova serrata*. A voucher specimen was deposited in the National Museum of Nature and Science (TNS), Tsukuba, Japan. A living ex-type culture was deposited in the Biological Resource Center of the National Institute for Technology and Evaluation (NBRC). *Cylindromonium lichenicola* strains (CBS 188.70 and CBS 415.70A) were also examined for comparison purposes.

Fungal Systematics and Evolution is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License



Morphological observations

Samples were observed using a dissecting microscope [M165 C (Leica, Wetzlar, Germany)] and a differential interference contrast compound microscope [BX53 (Olympus, Tokyo, Japan)]. Anatomical examination was performed using hand-cut sections mounted in a drop of water or clear lactophenol. Photographs were taken using a microscope digital camera [Flexacam C3 (Leica, Wetzlar, Germany) or DP23 (Olympus, Tokyo, Japan)]. Dimensions of ascospores, conidia, conidial mass, phialide and hyphal width are given as (minimum–) range of mean ± standard deviation (–maximum) (n = number of measurements). Chemical reactions of the perithecia were observed by using 10 % KOH. To determine if there is a significant difference between each dimension of the present fungus and *C. lichenicola*, the t-test was performed using Microsoft Excel.

Isolation of fungal cultures

Fungal cultures were isolated from freshly collected material. Mycelium or single ascospores were picked up using a flamed needle and plated on 1 % malt extract agar (MEA). To confirm differences in colony characteristics on each agar medium, mycelial plugs were subcultured on 1 % MEA, potato dextrose agar (PDA) (Nissui Pharmaceutical, Tokyo, Japan), oatmeal agar (OA) (Becton Dickinson and Co, New Jersey, USA), Sabouraud maltose agar (SMA) (Thermo Fisher Scientific, Massachusetts, USA), malt yeast extract agar (MYA) (Ahmadjian 1961) and Sabouraud glucose agar (SGA) (Stocker-Wörgötter 2002), confirming the recipes of these media according to Crous *et al.* (2019a). Colour of colonies were determined based on Kornerup & Wanscher (1978).

DNA extraction, PCR amplification and sequencing

Perithecia were sampled from specimen TNS-L-131533, mycelium from specimen TNS-L-131534, and mycelia from a culture derived from specimen TNS-L-131535. For DNA extraction, fungal tissues were suspended in 20 μ L of DNA extraction buffer [10 mM Tris-HCl (pH 8.3), 1.5 mM MgCl₂, 50 mM KCl, 0.01 % sodium dodecyl sulfate (SDS), 0.01 % Proteinase K], incubated at 37 °C for 60 min, and denatured 90 °C for 10 min; 30 μ L of sterile distilled water (SDW) added to the tubes, and stored in a freezer at -20 °C.

Partial sequences of the nuc rDNA ITS1-5.8S-ITS2 (ITS), large subunit (LSU) nuc rDNA regions, elongation factor 1-alpha (tef1), and RNA polymerase II second largest subunit (rpb2) were amplified as these regions are frequently used for phylogenetic analyses of Nectriaceae (Lombard et al. 2015, Crous et al. 2019b, 2020, 2021). The ITS region was amplified using the primers ITS5 and ITS4 (White et al. 1990), LSU rDNA using primers LIC24 (Miadlikowska & Lutzoni 2000) and LR7 (Vilgalys & Hester 1990) or LROR (Rehner & Samuels 1994) and LR6 (Vilgalys & Hester 1990), tef1 using EF1-983F and EF1-1567R (Rehner & Buckley 2005), and rpb2 using RPB2-5F2 and RPB2-7cR (O'Donnell et al. 2007). PCR was performed in a 15 µL reaction volume containing 1 µL DNA template, 7.5 µL GenRED PCR Mix Plus (Nippon Gene, Tokyo, Japan), 1.5 μL each primer (2 pmol/μL), and 3.5 μL distilled water. The PCR was performed in a TaKaRa PCR Thermal Cycler Dice® Touch (TaKaRa, Shiga, Japan) as follows for the ITS region; 5 min at 95 °C, followed by 40 cycles of 30 s at 94 °C, 30 s at 53 °C, 1 min at 72 °C, and a final step of 8 min at 72 °C. PCR conditions for LSU, *tef1* and *rpb2* were set according to Frisch *et al.* (2020), Rehner & Buckley (2005) and O'Donnell *et al.* (2007), respectively.

PCR products were checked by electrophoresis on a 1.5 % agarose gel stained with Midori Green Direct DNA Stain (Nippon Genetics, Tokyo, Japan) and visualised using WSE-5200 Printgraph 2 M (ATTO, Tokyo, Japan). The PCR products were purified using a FastGene[™] Gel/PCR extraction kit (Nippon Genetics, Tokyo, Japan) and ExoSAP-IT[™] (Thermo Fisher Scientific, Massachusetts, USA) following the manufacturer's instructions.

Sequences were obtained via a DNA sequencing service using Applied Biosystems 3730xl DNA analyzer (Thermo Fisher Scientific, Massachusetts, USA) (Eurofins Genomics, Tokyo, Japan). Data and accession number of the voucher specimen, and the obtained sequences from the International Nucleotide Sequence Database (INSD) are shown in Table 1.

Phylogenetic analysis and comparison of sequence data

Both newly generated ITS sequences and reference sequences were used in the phylogenetic analysis. Sequence data of loci other than ITS do not exist for some isolates of the genus *Cylindromonium*. Therefore, only ITS, LSU and *rpb2* of four *Cylindromonium* species were used for the phylogenetic analysis (Table 1). ITS sequence data of species with relatively high identity (> 85 %) in the BLAST search and other hypocrealean fungi in the *Nectriaceae* were included in the analysis to infer the taxonomic position of the targeted fungi (Table 1). In these analyses, *Stachybotrys chartarum* (KM231858) was chosen as the outgroup (Lombard *et al.* 2015). Sequences of each locus (ITS, LSU, *tef1* and *rpb2*) were compared with those of *C. lichenicola*. All sequences analysed in this study were deposited in the DNA Data Bank of Japan (DDBJ), a member of International Nucleotide Sequence Database Collaboration (INSDC).

Assembling forward and reverse strands of the sequenced loci were carried out with MUSCLE v. 3.6 (Edgar 2004) in MEGA v. 7 (Kumar *et al.* 2016) to obtain consensus sequences. DNA sequences were aligned using the online version MAFFT v. 7 (Katoh *et al.* 2019) (https://mafft.cbrc.jp/alignment/server/) with default settings. MEGA v. 7 (Kumar *et al.* 2016) was used to truncate sequences up to the determined edge of the dataset.

Phylogenetic analyses were performed with Maximum likelihood (ML) using an online version W-IQ-Tree v. 1.6.12 (Trifinopoulos *et al.* 2016) (http://iqtree.cibiv.univie.ac.at/). All characters were equally weighted, and gaps were treated as missing data. The ML analysis for the ITS region alignment using the TIM2+F+I+G4 model and for a combined alignment of the three loci, ITS, LSU, *rpb2* using the TN93+G (for ITS and LSU) and TN93+I (for *rpb2*) were performed with 1 000 bootstrap replicates. FigTree v. 1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/) and MEGA X were used for plotting the phylogenetic trees. Sequence alignments were deposited in TreeBASE (http://purl.org/phylo/treebase/phylows/study/TB2:S30024).

Inoculation experiments

Symptomless thalli of *D. applanata*, the host lichen of *C. dirinariae*, and those of the non-host lichen *Parmotrema tinctorum* were collected in Tsukuba city, Ibaraki prefecture, Japan and confirmed as non-infected via microscopy. Following this step, the lichen surface was cleaned using an ultrasonic cleaner with 0.005 % Aerosol® OT (a surface-active agent) for 1

Species	Strain numbers	Host/Substrate	Collection sites	ITS Accession No.	LSU Accession No.	<i>tef1</i> Accession No.	<i>rpb2</i> Accession No.	References
Calostilbe striispora	CBS 133491	Erythrina glauca	Trinidad and Tobago	KM231789	I	I	I	Lombard <i>et al</i> . (2015)
Ciliciopodium brevipes	CBS 691.83	Fagus sylvatica	Netherlands	KM231856	I	I	I	Lombard <i>et al.</i> (2015)
Ciliciopodium hyalinum	CBS 106.13	Soil	Switzerland	KM231857	I	I	I	Lombard <i>et al.</i> (2015)
Cylindrium aeruginosum	CBS 693.83	Fagus sylvatica	Netherlands	KM231854	I	I	I	Lombard <i>et al.</i> (2015)
Cylindrium elongatum	CBS 685.83A	Fagus sp.	Netherlands	KM231852	I	I	I	Lombard <i>et al.</i> (2015)
Cylindromonium alloxyli	CPC 38159	Meliola on leaves of Alloxylon pinnatum	Austria	MW175339	MW175379	I	MW173114	Crous <i>et al.</i> (2020)
Cylindromonium eugeniicola	CPC 37170	Leaf litter of <i>Eugenia capensis</i>	South Africa	NR166338	I	I	I	Crous <i>et al</i> . (2019b)
Cylindromonium everniae	CBS 148255	Evernia prunasti	Netherlands	NR175231	OK664736	I	NR175231	Crous <i>et al</i> . (2021)
Cylindromonium dirinariae sp. nov.	TNS-L-131533	Dirinaria applanata	Japan	LC731273	LC731274	LC731275	LC744391	This study
	TNS-L-131534	Dirinaria applanata	Japan	LC731276	LC744402	LC744396	LC744390	This study
	TNS-L-131535	Dirinaria applanata	Japan	LC731277	LC744401	LC744395	LC744392	This study
Cylindromonium lichenicola	CBS 415.70A	Aerial algae	Netherlands	MH859774	LC744400	LC744397	LC744394	Vu <i>et al.</i> (2019); This study
	CBS 303.70	Alnus sp.	Germany	MH859675	I	I	I	Vu <i>et al.</i> (2019)
	CBS 188.70	Apothecia of lichen	Germany	MH859549	LC744399	LC744398	LC744393	Vu <i>et al.</i> (2019); This study
Cylindromonium rhabdosporum	CBS 438.66	Cladonia furcata	Austria	MH858850	I	I	I	Vu <i>et al.</i> (2019)
Falcocladium multivesiculatum	CBS 120386	Leaf litter	Brazil	JF831936	I	I	I	Rungjindamai <i>et al.</i> (unpublished)
Falcocladium sphaeropedunculatum	CBS 111292	Leaf litter	Brazil	JF831938	I	I	I	Rungjindamai <i>et al.</i> (unpublished)
Falcocladium thailandicum	CBS 121717	Eucalyptus camaldulensis	Thailand	JF831939	I	I	I	Rungjindamai <i>et al.</i> (unpublished)
Hyaloseta nolinae	CBS 109837	Nolina micrantha	NSA	KM231846	I	I	I	Lombard <i>et al.</i> (2015)
Lectera colletotrichoides	CBS 109728	Medicago sativa	Turkey	KM231851	I	I	I	Lombard <i>et al.</i> (2015)
Nectria balansae	CBS 123351	<i>Coronila</i> sp.	France	HM484552	I	I	I	Hirooka <i>et al.</i> (2011)
Nectria cinnabarina	CBS 125165	Aesculus sp.	France	HM484548	I	I	I	Hirooka <i>et al</i> . (2011)
Nectria dacryocarpa	CBS 121.87	Tree fern	Sulawesi	KM231850	I	I	I	Lombard <i>et al.</i> (2015)
Nectria mariae	CBS 125294	Buxus sempervirens	France	JF832629	I	I	I	Hirooka <i>et al.</i> (2012)
Nectria nigrescens	CBS 125148	Wood	NSA	HM484707	I	I	I	Hirooka <i>et al.</i> (2011)
Phialoseptomonium eucalypti	CBS 145542	Leaves of <i>Eucalyptus</i>	Australia	MK876402	I	I	I	Crous <i>et al.</i> (2020)
Pochonia sp.	CBS 634.75	Arcyria sp.	Netherlands	KM231845	I	I	I	Lombard <i>et al.</i> (2015)
	CBS 892.7	Myxomycete	Netherlands	KM231844	I	I	I	Lombard <i>et al.</i> (2015)
	CBS 401.70	Myxomycete	Netherlands	KM231843	I	I	I	Lombard <i>et al.</i> (2015)
Rodentomyces reticulatus	CBS 128675	Rodent dung	Italy	JF832659	I	I	I	Hirooka <i>et al.</i> (2012)

Table 1. Sources of DNA sequence data used in phylogenetic analyses and comparison of sequence data.

3

Table 1. (Continued).								
Species	Strain numbers	Host/Substrate	Collection sites	ITS Accession No.	LSU Accession No.	<i>tef1</i> Accession No.	<i>rpb2</i> Accession No.	References
Sarocladium kiliense	CBS 400.52	Ficus carica	UK	KM231849	I	I	I	Lombard <i>et al.</i> (2015)
Septofusidium berolinense	CBS 731.70	1	Germany	KM231841	I	I	I	Lombard <i>et al.</i> (2015)
Stachybotrys chartarum	CBS 129.13	I	Unknown	KM231858	MH866145	I	KM232434	Lombard <i>et al.</i> (2015)
Thyronectria lamyi	CBS 417.89	Berberis vulgaris	Germany	KM231837	Ι	I	I	Lombard <i>et al.</i> (2015)
Thyronectria pyrrhochlora	CBS 125131	Acer campestre	Austria	HM484545	I	I	I	Hirooka <i>et al</i> . (2011)
Thyronectria quercicola	CBS 128976	Quercus ilex	Spain	JF832624	I	I	I	Hirooka <i>et al</i> . (2012)
Thyronectria sinopica	CBS 462.83	Hedera helix	Netherlands	HM484542	Ι	I	I	Hirooka <i>et al</i> . (2011)
Tilachlidium brachiatum	CBS 505.67	Hypholoma fasciculare	Poland	KM231839	I	I	I	Lombard <i>et al.</i> (2015)
Trichonectria rectipila	CBS 132.87	I	USA	MH862058	I	I	I	Vu <i>et al.</i> (2019)
Trichonectria setadpressa	J.E.20–13	Lobariella pallida	France	MT153969	I	I	I	Flakus <i>et al.</i> (2019)
CBS: Westerdiik Eungal Biodiversity In	stitute Litrecht Net	-herlands. CPC. Culture collection	of Pedro Crois	housed at the V	Vecterdiik Eungal	Rindiversity Institu	ute. I E · The nrivat	e herharium of lavier Ftavo

min. This step was repeated five times. After this process, samples were rinsed with distilled water. Agar pieces including hyphae of isolates of *C. dirinariae* and *C. lichenicola* were inoculated onto the lichen thalli and maintained in 90-mm-diam glass Petri dishes with a sheet of wet filter paper to retain a high humidity at room temperature. Lichens inoculated were misted from above and moistened when they began to dry, every 2 to 3 d. In addition, some lichens were not inoculated and maintained under the above conditions as a control.

RESULTS

Taxonomy

Classification: Nectriaceae, Hypocreales, Sordariomycetes.

Cylindromonium dirinariae Ohmaki & Okane, *sp. nov.* MycoBank MB 846061. Fig. 1.

Etymology: Name refers to the host genus *Dirinaria* from which it was isolated.

Diagnosis: Ascomata perithecial, globose, orange; *phialides* short; *conidia* aggregated in mucoid packets in the apex of phialides, ellipsoid, aseptate; the species differs from all other *Cylindromonium* species by its characteristic DNA sequences (ITS, LSU, *tef1*, *rpb2*) and from its closest relative *C. lichenicola* also by its shorter phialides.

Description: Ascomata occur on the upper surface or soralia of the host lichen thallus; perithecial, scattered, globose, 80-100 µm diam, poculiform when dry, pale orange when young but later becoming dark, KOH negative, ascomatal wall layers of textura globosa, 14-18 µm thick. Asci broadly cylindrical to clavate, non-stipitate, (21.2–)24.5 \pm 2.3(-29.1) × (3.9-)5.0 \pm 0.8(-6.5) µm (n = 12), unitunicate, apex simple, 8-spored. Ascospores biseriate, ellipsoid, hyaline, smooth, medially 1-septate, $(5.0-)7.8 \pm 1.1(-10.1) \times (1.8-)2.6 \pm 0.4(-3.7)$ μ m, length/breadth (l/b) = (2.1–)2.6 ± 0.4(–3.9) (n = 40). Mycelium consisting of hyaline, smooth, septate, branched, 2 µm diam hyphae. Conidiogenous cells arising directly from aerial hyphae, hyaline, smooth, subcylindrical, 20–35 µm tall, 2 µm wide at the base, tapering to 1 µm at the apex, phialidic, with non-flared collarette. Conidia solitary, adhering in a slimy mass, hyaline, smooth, aseptate, ellipsoid with obtuse ends, $(4.1-)5.9 \pm 0.9(-10.3) \times (1.5-)2.4 \pm 0.4(-3.5) \mu m$, I/b $= (1.5-)2.55 \pm 0.5(-3.9) (n = 75).$

Culture characteristics: Colonies flat, circular or irregular, with moderate aerial mycelium and smooth, lobate margin, reaching 20 mm diam on MEA, PDA, OA, SMA and MYA, 30 mm diam on SGA after 3 wk at 23 °C in darkness. On MEA, PDA, SMA, SGA and MYA surface and reverse strong brownish orange; on OA surface grayish orange.

Host: Dirinaria applanata.

Distribution: Japan.

Typus: **Japan**, Ibaraki, Tsukuba, Tennodai, Univ. of Tsukuba, 36°06′08″N, 140°06′24″E, lichenicolous on *Dirinaria applanata* on bark of *Zelkova serrata*, 1 Aug. 2020, *A. Ohmaki, I. Okane, K. Ohmachi, K. Miyazawa & K. Gibu*, FAO 005 (**holotype** TNS-L-131533, culture ex-type NBRC 115852); DDBJ: ITS = LC731273; LSU = LC731274; *tef1* = LC731275; *rpb2* = LC744391.

Pamplona





Fig. 1. *Cylindromonium dirinariae*. **A.** *C. dirinariae* colonizing on *Dirinaria applanata*. **B, C.** Colonies on MEA. B. Surface. C. Reverse. **D.** Growth habit on *Dirinaria applanata*. **E.** Perithecium. **F.** Conidiogenous cells and conidia. **G.** Asci. **H.** Ascospores. **I.** Conidia. **J.** Phialide and conidial mass. Scale bars: B, C = 3 cm; D = 0.5 mm; E, F = 0.25 mm; G, I, J = 10 µm; H = 5 µm.



Additional materials examined: Japan, Ibaraki, Tsukuba, Tennodai, Univ. of Tsukuba, 36°06′08″N, 140°06′24″E, from *Dirinaria applanata* on the bark of *Zelkova serrata*, 1 Aug. 2020, *A. Ohmaki, I. Okane, K. Ohmachi, K. Miyazawa & K. Gibu*, NBRC 115851 = TNS-L-131534 (FAO 004), DDBJ: ITS = LC731276; LSU = LC744402; *tef1* =LC744396; *rpb2* =LC744390; 6 Nov. 2020, *A. Ohmaki*, TNS-L-131536 (FAO 090), in an inoculation experiment with isolates on *D. applanata* (Ibaraki: Tsukuba, 1 Aug. 2020, *A. Ohmaki, I. Okane, K. Ohmachi, K. Miyazawa & K. Gibu*, TNS-L-131533); Tsukuba, Tennodai, Univ. of Tsukuba, 36°06′38″N, 140°06′16″E, from *Dirinaria applanata* on the bark of *Zelkova serrata*, 15 Mar. 2021, *A. Ohmaki*, NBRC 115853 = TNS-L-131535 (FAO 006), DDBJ: ITS = LC731277; LSU = LC744401; *tef1* = LC744395; *rpb2* = LC744392.

Notes: The ex-type strain of *C. lichenicola* (CBS 425.66) was not available for study, so we conducted morphological observations on other available strains of *C. lichenicola* (CBS 188.70 and CBS 415.70A) that are similar in terms of their collection sites and hosts. In addition, these strains were also studied when the genus *Cylindromonium* was established in Crous *et al.* (2019b).

Morphological features of the *C. lichenicola* strains (CBS 188.70 and CBS 415.70A) shown in Table 2 correlated well with the original description provided by Gams (1971) [conidia size $5.5-9.8 \times 1.5-2.5 \ \mu m \ (l/b = 3.0-4.4)$, phialide length $30-60 \ \mu m$, phialide width base $2.0-3.0 \ \mu m$, apex $0.7-1.5 \ \mu m$].

As a result of the t-test, there were significant differences between *C. dirinariae* and *C. lichenicola* in conidia, phialide length and width (Table 2). As for phialide length, phialides of *C. dirinariae* were about half as long as those of *C. lichenicola*.

Materials examined of *Cylindromonium lichenicola*: **Germany**, Probsteierhagen, Schüttbrehm, from unnamed apothecia of lichen on tree bark, Oct. 1996 (living strain CBS 188.70), GenBank: LSU = LC744399; *tef1* = LC744398; *rpb2* = LC744393. **Netherlands**, Utrecht, Amelisweerd, from aerial algae on tree bark, Oct. 1968 (living strain CBS 415.70A), GenBank: LSU = LC744400; *tef1* = LC744397; *rpb2* = LC744394.

Phylogeny

The ITS sequences derived from DNA extracted from a perithecium (specimen TNS-L-131533) and that from mycelia from another specimen (TNS-L-131534) of *C. dirinariae* were identical. The Phylogenetic analysis based on the ITS region revealed that all three sequences, adding that from mycelia (TNS-L-131535) to the above two, were grouped in a fully supported clade (Fig. 2). The clade was sister to *C. lichenicola* and linked with five *Cylindromonium* species including the type species of the genus *Cylindromonium*, *C. everniae*. The clade including the *Cylindromonium* species also included species of *Trichonectria* (*Bionectriaceae*) and *Phialoseptomonium* (*Nectriaceae*).

Sequences of LSU, *tef1*, and *rpb2* derived from specimen TNS-L-131533, TNS-L-131534 and TNS-L-131535 of *C. dirinariae* were identical. The ITS sequence from TNS-L-131535 differed

from the other two sequences from TNS-L-131533 and TNS-L-131534 in one site. While the sequences of ITS, LSU, and *tef1* of *C. lichenicola* strains CBS 188.70 and CBS 415.70A were identical, those of *rpb2* of the two stains were different in 18 sites. Comparison of *C. dirinariae* with *C. lichenicola* (CBS 188.70 and CBS 415.70A) showed that there were 19 gap sites (96.7 % in nucleotide identity) in ITS, 11 gap sites including 2 bp deletions (98.8 %) in LSU, 26 gap sites in *tef1* (93.7 %), 153 gap sites (CBS 415.70A *vs. C. dirinariae*) or 155 gap sites (CBS 188.70 *vs C. dirinariae*) including 44 bp insertions in *rpb2* (84.9–85.1 %). Phylogenetic analysis based on the concatenated sequences of ITS, LSU and *rpb2* showed that three sequences of *C. dirinariae* grouped together with full bootstrap support and clearly segregated from *C. lichenicola* and other species (Fig. 3).

Inoculation experiments

The two *Cylindromonium* spp. studied were able to colonise the inoculated lichens, except for failures due to contamination, extreme dryness or moisture (Fig. 4; Table 3).

Cylindromonium dirinariae colonised and produced perithecia on the thalli of *D. applanata*. Colonies reached 5 mm diam about 1 wk post inoculation. Pinkish discolouration was observed in the part of lichen's thalli covered with hyphae of *C. dirinariae*. The asexual morph also developed on inoculated thalli. Perithecia developed 2 wk post inoculation. Morphological features of *C. dirinariae* on the inoculated lichens coincided well with those of *C. dirinariae* observed in the field.

On the other hand, although *C. lichenicola* also colonised and produced the asexual morph on *D. applanata*, no perithecia were produced.

Cylindromonium dirinariae and *C. lichenicola* colonised and sporulated asexually, but no perithecia were produced on *P. tinctorum*. The lichen's thalli were covered with hyphae of the lichenicolous fungi, and discoloured brownish around the point of inoculation. Colonies reached 5 mm diam after 1 wk and 1.5–2 cm diam after 2 wk post inoculation. Lichens used for controls remained healthy. In addition, three single ascospore cultures were obtained of *C. dirinariae*, and each culture was inoculated onto thalli of *D. applanata*. As a result, they colonised the lichen and produced perithecia.

DISCUSSION

In the molecular phylogenetic analysis based on ITS sequence data and concatenated sequences of three loci, the three sequences of *C. dirinariae* clustered in a single clade and positioned as sister to *C. lichenicola*, which is the most closely related species in morphology and DNA phylogeny (identity = 96.7 %), supported by high bootstrap values. Furthermore, *C. dirinariae* was related to the group consisting of *C. everniae* and

Table 2. Comparison of dimensions (µm) between Cylindromonium dirinariae and C. lichenicola.

		,			
	Conidia*	Phialide length*	Phialide width (Base)*	Phialide width (Apex)*	Hyphae width
C. dirinariae	(4.1–)5.9 ± 0.9(–10.3) × (1.5–)2.4 ± 0.4(–3.5)	(14.7–)27.4 ± 8.1(– 56.7)	(1.7–)2.3 ± 0.3(–3.1)	(1.0–)1.4 ± 0.2(–2.0)	(1.2–)2.0 ± 0.3(–2.6)
C. lichenicola	(4.1–)6.8 ± 1.1(–10.0) × (1.5–)2.4 ± 0.5(–3.3)	(33.9–)51.7 ± 7.6(– 69.9)	(2.0–)3.2 ± 0.5(–4.0)	(1.1–)1.6 ± 0.3(–2.5)	(1.5–)2.3 ± 0.5(–3.3)
*Significantly	lifferent				

*Significantly different.





Fig. 2. A phylogenetic tree of *Cylindromonium dirinariae* and other *Nectriaceae* and hypocrealean fungi constructed from a maximum-likelihood (ML) analysis based on the ITS sequences. The outgroup is *Stachybotrys chartarum*. Number at the nodes represents the bootstrap value.

C. rhabdosporum, supported by high bootstrap values (99 %) (Fig. 2). Comparison of other loci sequences also showed that similarity between *C. dirinariae* and *C. lichenicola* were 96.7 % in ITS, 98.8 % in LSU, 93.7 % in *tef1*, 84.9–85.1 % in *rpb2*. In addition to phylogeny, *C. dirinariae* is also morphologically distinct. In our morphological observations and the inoculation experiments, *C. dirinariae* had 20–35 µm tall phialides and produced perithecia on *D. applanata*, while *C. lichenicola* had longer phialides (45–60 µm tall), and failed to produce perithecia on *D. applanata* and *P. tinctorum*. Hence, we concluded that *C. dirinariae* represents

a new species. In addition, this is the first report of the sexual morph for the genus *Cylindromonium*.

The genus *Cylindromonium* has been reported from Belgium (Gams 1971, Diederich 1989), Germany (Gams 1971, Brackel 2010), Great Britain (Hitch 1995), France (Roux 2012), Luxembourg (Diederich 1989), the Netherlands (Brand *et al.* 2013, Crous *et al.* 2021), Czech Republic (Kocourková 2009), India (Joshi *et al.* 2016), Ukraine (Khodosovtsev *et al.* 2018), and Australia (Gams 1971, Crous *et al.* 2020). This is the first report of the genus *Cylindromonium* from Japan.

Table 3. Colonization rate in inoculation experiments.

		Inoculum	
Lichen species inoculated	Cylindromonium dirinariae	Cylindromonium lichenicola	Control
Dirinaria applanata	16/20* (80 %)**	13/15 (87 %)	3/12 (25 %)
Parmotrema tinctorum	4/4 (100 %)	2/2 (100 %)	0/5 (0 %)

*Number of colonised lichenicolous fungi/the number of inoculated lichens.

^{**}Perithecia produced.



Fig. 3. A phylogenetic tree for *Cylindromonium dirinariae* and other *Cylindromonium* spp. constructed from a maximum-likelihood (ML) analysis based on concatenated sequence dataset of ITS, LSU rDNA and *rpb2*. The outgroup is *Stachybotrys chartarum*. Number above a branch represents the bootstrap value.

Inoculation experiments using single ascospore cultures revealed that this fungus is homothallic. The newly described *C. dirinariae* was isolated from *D. applanata*, while the hosts of *C. lichenicola* are diverse, including the lichens *Cladonia*, *Hypogymnia*, *Parmelia saxatilis*, *Tremella cladoniae*, lichens overgrowing *Stereum* species, fungi; *Bulgaria inquinans*, algalcovered bark, *Alnus* bark and *Betula* litter (Gams 1971, Diederich 1989, Hitch 1995, Brackel 2010, Roux 2012, Brand *et al.* 2013, Tsurykau *et al.* 2016). Results of our inoculation experiments are suggestive of host specificity; *i.e.*, *C. dirinariae* was able to form perithecia on its original host lichen, *D. applanata*, but not on *P. tinctorum. Cylindromonium dirinariae* could therefore be considered host specific.

Glenn *et al.* (1997) found that *Cylindromonium rhabdosporum* (as *Acremonium rhabdosporum*), occurred on healthy-looking thalli in the field, and perithecia of *Nectriopsis rubefaciens* (as *Nectria rubefaciens*) appeared on the same thalli in the closed plates. Although they mentioned the relationship between *C. rhabdosporum* and *N. rubefaciens*, they did not confirm that they belong to the same holomorph. Further study is therefore needed to determine the homogeneity between *C. rhabdosporum* and *N. rubefaciens*.

Presently there are only a few reports of inoculation experiments of lichenicolous fungi, and they were conducted in the field (Fatma *et al.* 2019). Glenn *et al.* (1997) found that continuously moist conditions probably play a pivotal role for lichenicolous fungi to produce perithecia. Inoculation experiments in the moist condition in Petri dishes using axenic cultures of lichenicolous fungi may therefore be a useful technique for

studying morphological, physiological and ecological features of lichenicolous fungi. This work has demonstrated the potential of inoculation experiments to investigate the morphological feature of perithecia and host specificity.

Presently the ecology of *C. dirinariae* remains unclear. Other species of *Cylindromonium* were reported to be mycophilic or saprophytic (Crous *et al.* 2019b, 2020). We expect that inoculation experiments will reveal the interaction between lichenicolous fungi and their host lichens which could help us to better understand the ecological role of lichenicolous fungi.

The ascomycete family *Nectriaceae* includes numerous important plant and human pathogens as well as several facultatively fungicolous or insecticolous species (Rossman 1996, Lombard *et al.* 2015). Members of *Nectriaceae* are characterised by uniloculate ascomata that are white, yellow, orange-red or purple, unitunicate asci and phialidic asexual morphs (Rossman *et al.* 1999, Lombard *et al.* 2015). In many cases ascomata show a change of colour when mounted in KOH (Lombard *et al.* 2015). In our study, perithecia of the *C. dirinariae* were orange in colour and did not react in KOH.

About 400 genera and 2 300 species of lichenicolous fungi are known from lichens, but the actual number of lichenicolous fungal species could be much higher (Diederich *et al.* 2018). Diederich *et al.* (2018) estimated 3 000 – 5 000 lichenicolous fungal species will eventually be described based on Hawksworth's global estimates of fungal diversity (Hawksworth 1991, 2001) and the total number of lichen species (Lücking *et al.* 2017a, 2017b). Lichenicolous fungi are assumed to be an important source of new species in many groups of fungi, including *Nectriaceae*.





Fig. 4. Lesions on the lichen thalli after inoculation. A. Colony of *C. dirinariae* on *D. applanata*. B. Colony of *C. dirinariae* on *P. tinctorum* C. Colony of *C. lichenicola* on *D. applanata*. D. Colony of *C. lichenicola* on *P. tinctorum*. Scale bars = 5 mm.

ACKNOWLEDGEMENTS

This research was partially financially supported by Japan Society for the Promotion of Science (JSPS) KAKENHI Grant Number 21K19286 (IO).

Conflict of interest: The authors declare that there is no conflict of interest.

REFERENCES

- Ahmadjian V (1961). Studies on lichenized fungi. *The Bryologist* **64**: 168–179.
- Brackel WV (2010). Weitere Funde von flechtenbewohnenden Pilzen in Bayern. Beitrag zu einer Checkliste V. *Berichte der Bayerischen Botanischen Gesellschaft* **80**: 5–32.
- Brand AM, Sparrius LB, Aptroot A (2013). 31 nieuwe soorten korstmossen en lichenicole fungi voor Nederland. *Buxbaumiella* **97**: 17–22.
- Crous PW, Verkley GJM, Groenewald JZ, *et al.* (eds) (2019a). *Fungal Biodiversity*. [*Westerdijk Laboratory Manual Series No. 1.*]. Utrecht: Westerdijk Fungal Biodiversity Institute, Utrecht, the Netherlands.
- Crous PW, Wingfield MJ, Lombard L, *et al.* (2019b). Fungal Planet description sheets: 951–1041. *Persoonia* **43**: 223–425.
- Crous PW, Cowan DA, Maggs-Kölling G, et al. (2020). Fungal Planet description sheets: 1112–1181. Persoonia **45**: 251–409.

- Crous PW, Osieck ER, Jurjevi Ž, *et al.* (2021). Fungal Planet description sheets: 1284–1382. *Persoonia* **47**: 178–374.
- Diederich P (1989). Les lichens épiphytiques et leurs champignons lichénicoles (macrolichens exceptés) du Luxembourg. *Travaux Scientifiques du Musée National D'Histoire Naturelle de Luxembourg* 14: 1–268.
- Diederich P, Lawrey JD, Ertz D (2018). The 2018 classification and checklist of lichenicolous fungi, with 2000 nonlichenized, obligately lichenicolous taxa. *The Bryologist* **121**: 340–425.
- Edgar RC (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* **32**: 1792–1797.
- Fatma MAEW, Abdou MMM, Gehad MEH, et al. (2019). Efficiency of lichenicolous fungi in controlling citrus lichen Xanthoria parietina. Annals of Agricultural Science Moshtohor 57: 507–516.
- Flakus A, Etayo J, Miadlikowska J, *et al.* (2019). Biodiversity assessment of ascomycetes inhabiting *Lobariella* lichens in Andean cloud forests led to one new family, three new genera and 13 new species of lichenicolous fungi. *Plant and Fungal Systematics* **64**: 283–344.
- Frisch A, Grube M, Kashiwadani H, et al. (2018). Arthoniaceae with reddish, K+ purple ascomata in Japan. Phytotaxa **356**: 19–33.
- Frisch A, Tadome K, Moon KH, et al. (2020). Phylogenetic status of Arthonia phaeophysciae (Arthoniaceae, Ascomycota), a species new to Japan. The Journal of Japanese Botany 95: 133–140.
- Gams W (1971). Cephalosporium-artige Schimmelpilze (Hyphomycetes). Gustav Fischer Verlag, Stuttgart.



- Glenn MG, Gomez-Bolea A, Orsi EV (1997). Effects of thallus damage on interactions of lichens with non-lichenized fungi under natural and laboratory conditions. *The Lichenologist* **29**: 51–65.
- Hawksworth DL (1991). The fungal dimension of biodiversity: magnitude, significance, and conservation. *Mycological Research* **95**: 641–655.
- Hawksworth DL (2001). The magnitude of fungal diversity: the 1.5 million species estimate revisited. *Mycological Research* **105**: 1422–1432.
- Hirooka Y, Rossman AY, Chaverri P (2011). A morphological and phylogenetic revision of the *Nectria cinnabarina* species complex. *Studies in Mycology* 68: 35–56
- Hirooka Y, Rossman AY, Samuels GJ, et al. (2012). A monograph of Allantonectria, Nectria, and Pleonectria (Nectriaceae, Hypocreales, Ascomycota) and their pycnidial, sporodochial, and synnematous anamorphs. Studies in Mycology **71**: 1–210.
- Hitch C (1995). New, rare and interesting British lichen records. *British Lichen Society Bulletin* **76**: 47–59.
- Joshi Y, Falswal A, Tripathi M, *et al.* (2016). One hundred and five species of lichenicolous biota from India: An updated checklist for the country. *Mycosphere* **7**: 268–294.
- Katoh K, Rozewicki J, Yamada KD (2019). MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* **20**: 1160–1166.
- Khodosovtsev A, Darmostuk V, Suija A, *et al.* (2018). *Didymocyrtis trassii sp. nov.* and other lichenicolous fungi on *Cetraria aculeata*. *The Lichenologist* **50**: 529–540.
- Kocourková J (2009). Observations on the genus *Neolamya*, with the description of the new species *N. xanthoparmeliae* (*Ascomycota*, genera *incertae sedis*). *Opuscula Philolichenum* **6**: 137–148.
- Kornerup A, Wanscher JH (1978). *Methuen handbook of colour, 3rd edition*. 1–252. Eyre Methuen, London.
- Kumar S, Stecher G, Tamura K (2016). MEGA7: molecular evolutionary genetics analysis version 70 for bigger datasets. *Molecular Biology* and Evolution **33**: 1870–1874.
- Lombard L, Van der Merwe NA, Groenewald JZ, *et al.* (2015). Generic concepts in *Nectriaceae*. *Studies in Mycology* **80**: 189–245.
- Lücking R, Hodkinson BP, Leavitt SD (2017a). The 2016 classification of lichenized fungi in the *Ascomycota* and *Basidiomycota* Approaching one thousand genera. *The Bryologist* **119**: 361–416.
- Lücking R, Hodkinson BP, Leavitt SD (2017b). Corrections and amendments to the 2016 classification of lichenized fungi in the *Ascomycota* and *Basidiomycota*. *The Bryologist* **120**: 58–69.
- Miadlikowska J, Lutzoni F (2000). Phylogenetic revision of the genus *Peltigera* (lichen-forming *Ascomycota*) based on morphological, chemical, and large subunit nuclear ribosomal DNA data. *International Journal of Plant Sciences* **161**: 925–958.
- O'Donnell K, Sarver BA, Brandt M, *et al.* (2007). Phylogenetic diversity and microsphere array-based genotyping of human pathogenic fusaria, including isolates from the multistate contact lensassociated US keratitis outbreaks of 2005 and 2006. *Journal of Clinical Microbiology* **45**: 2235–2248.
- Ohmura Y, Kashiwadani H (2018). *Checklist of lichens and allied fungi of Japan*. National Museum of Nature and Science Tokyo, Tokyo.
- Rambaut A (2018). FigTree v1.4.4 [accessed 25 October 2020] http:// tree.bio.ed.ac.uk/software/figtree/
- Rehner SA, Buckley E (2005). A *Beauveria* phylogeny inferred from nuclear ITS and EF1-α sequences: evidence for cryptic diversification and links to *Cordyceps* teleomorphs. *Mycologia* **97**: 84–98.
- Rehner SA, Samuels GJ (1994). Taxonomy and phylogeny of *Gliocladium* analysed from nuclear large subunit ribosomal DNA sequences. *Mycological Research* **98**(6): 625–634.

- Roux C (2012). Liste des lichens et champignons lichénicoles de France. Bulletin de la Société Linnéenne de Provence **16**: 1–220.
- Rossman AY (1996). Morphological and molecular perspectives on systematics of the *Hypocreales*. *Mycologia* **88**: 1–19.
- Rossman AY, Samuels GJ, Rogerson CT, et al. (1999). Genera of Bionectriaceae Hypocreaceae and Nectriaceae (Hypocreales, Ascomycetes). Studies in Mycology 42: 1–248.
- Stocker-Wörgötter E (2002). Resynthesis of photosymbiodemes. In: *Protocols in lichenology* (Kranner I, Beckett RP, Varma AK eds). Springer, Berlin: 47-60.
- Summerbell RC, Gueidan C, Schroers HJ, *et al.* (2011). *Acremonium* phylogenetic overview and revision of *Gliomastix*, *Sarocladium*, and *Trichothecium*. *Studies in Mycology* **68**: 139–162.
- Tadome K, Ohmura Y (2021). Two lichenicolous fungi, Illosporium carneum and Ovicuculispora parmeliae (Bionectriaceae, Ascomycota), new to Japan. Bulletin of the National Museum of Nature and Science, Series B (Botany) 47: 21–25.
- Tadome K, Ohmura Y (2022). Six species of montane and subalpine lichenicolous fungi new to Japan. *The Journal of Japanese Botany* 97: 23–32.
- Tadome K, Ohmura Y, Frisch A (2018). Arthonia molendoi (Arthoniaceae, Ascomycota), a lichenicolous fungus, new to Japan. The Journal of Japanese Botany 93: 407–409.
- Tadome K, Ohmura Y, Chaki M (2022). Spirographa pyramidalis (Spirographaceae, Ascomycota), a lichenicolous fungus, new to Japan. The Journal of Japanese Botany **97**: 212–215.
- Trifinopoulos J, Nguyen LT, von Haeseler A, et al. (2016). W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Research* **44**: W232–W235.
- Tsurykau A, Suija A, Heuchert B, et al. (2016). New or otherwise interesting records of lichens and lichenicolous fungi from Belarus II. Herzogia 29: 164–175.
- Vilgalys R, Hester M (1990). Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* **172**: 4238–4246.
- Vu D, Groenewald M, De Vries M, *et al.* (2018). Large-scale generation and analysis of filamentous fungal DNA barcodes boosts coverage for kingdom fungi and reveals thresholds for fungal species and higher taxon delimitation. *Studies in Mycology* **91**: 23–36.
- White TJ, Bruns T, Lee S, et al. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR protocols: a guide to methods and applications* (Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds). Academic Press, San Diego (California): 315–322.
- Zhurbenko MP, Ohmura Y (2018a). Contributions to the knowledge of lichenicolous fungi on *Thamnolia*. *Opuscula Philolichenum* **17**: 368–373.
- Zhurbenko MP, Ohmura Y (2018b). *Perigrapha cetrariae*, a new lichenicolous ascomycete on *Cetraria* from Japan. *Folia Cryptogamica Estonica* **55**: 17–19.
- Zhurbenko MP, Ohmura Y (2019). New and interesting records of lichenicolous fungi from the TNS herbarium: Part I. *Opuscula Philolichenum* **18**: 74–89.
- Zhurbenko MP, Ohmura Y (2020). Contributions to the knowledge of lichenicolous fungi growing on baeomycetoid lichens and *Icmadophila*, with a key to the species. *Lichenologist* **52**: 437–453.
- Zhurbenko MP, Tadome K, Ohmura Y (2018). *Pronectria japonica sp. nov.* and a key to the lichenicolous fungi and lichens growing on *Ochrolechia. Herzogia* **31**: 494–504.